

Accretion Disk Instabilities, CDM models and their role in Quasar Evolution

Evanthia Hatziminaoglou^{1,2}, Aneta Siemiginowska² and Martin Elvis²
 eva@ast.obs-mip.fr, asiemiginowska@head-cfa.harvard.edu, elvis@head-cfa.harvard.edu

ABSTRACT

We have developed a consistent analytical model to describe the observed evolution of the quasar luminosity function. Our model combines black hole mass distributions based on the Press - Schechter theory of the structure formation in the Universe with quasar luminosity functions resulting from a physics-based emission model that takes into account the time-dependent phenomena occurring in the accretion disks. Quasar evolution and CDM models are mutually constraining, therefore our model gives an estimation of the exponent, n , of the power spectrum, $P(k)$, which is found to be $-1.8 \leq n \leq -1.6$. We were able to reject a generally assumed hypothesis of a constant ratio between Dark Matter Halo and the Black Hole mass, since the observed data could not be fitted under this assumption. We found that the relation between the Dark Matter Halos and Black Hole masses is better described by $M_{BH} = M_{DMH}^{0.668}$. This model provides a reasonable fit to the observed quasar luminosity function at redshifts higher than ~ 2.0 . We suggest that the disagreement at lower redshift is due to mergers. Based on the agreement at high redshift, we estimated the merger rate at lower redshift, and argue that this rate should depend on the redshift, like $(1+z)^3$.

Subject headings: accretion: accretion disks - cosmology:theory - quasars:general

1. Introduction

Since their discovery by Schmidt in the early 60's, quasars have been an intriguing phenomenon and the mystery surrounding them is still far from being resolved. Soon after being identified as galactic nuclei, accretion onto a supermassive black hole became the leading model for powering these objects (Lynden Bell, 1969), whose luminosities can far exceed those of galaxies, but whose space density is around two orders of magnitude lower than that of bright galaxies. The question of whether quasar activity was a short-lived phenomenon that occurred repeatedly during a galaxy's lifetime (Cavaliere & Padovani, 1989; Siemiginowska & Elvis, 1997), or whether quasars never "wake up" again after a single, longer-term,

activity cycle (Haiman & Loeb, 1998; Haehnelt, Natarajan & Rees, 1998), leaving behind massive black hole remnants, has not yet been resolved.

The quasar population clearly evolves with cosmic time. The quasar space density above a given luminosity increases by a factor of ~ 50 from the present to $z \sim 2.5$, then stalls and probably decreases to $z \sim 5$ (Hook, Shaver & McMahon, 1998). For many years the observed quasar evolution has been addressed phenomenologically, mostly via Pure Luminosity Evolution (PLE - Mathez, 1976) or Pure Density Evolution (PDE - Schmidt, 1968). Both descriptions have just one free adjustable parameter, but are physically rather abstract. Moreover PDE and PLE are observationally indistinguishable when only a power law is observed, as is common. Instead physically based models may be constrained by a simple power law, since neither the slope nor the normalization is now arbitrary. An early attempt to explain the observed quasar space den-

¹Laboratoire d'Astrophysique de Toulouse, Toulouse, France

²Harvard-Smithsonian Center for Astrophysics, Cambridge, USA

sity using a physics-based model was made by Efsthathiou and Rees (1988), who used the Press - Schechter formalism (Press & Schechter, 1974) for dark matter halo formation in order to predict the space density of the more luminous quasars. This work has been followed by several such attempts, with results that do not always agree (Cavaliere & Padovani, 1988; Haehnelt & Rees, 1993; Yi, 1996; Siemiginowska & Elvis, 1997; Haiman & Menou, 2000; Haehnelt, Natarajan & Rees, 1998; Salucci et al., 1999). Cavaliere & Vittorini (1998) suggest that the steep rise of the number density of quasars from early epochs to a redshift of ~ 2.5 is due to the newly formed galactic halos (density evolution) while their fall at lower redshifts is due to merging processes (luminosity evolution), creating a link between the phenomenological models and the more robust theory. Quasars are now a common ingredient of N-body simulations, semi-analytical or analytical models, which follow step by step the formation and evolution of galaxies within Dark Matter Halos and the merging processes (e.g. Richstone et al., 1998; Kauffmann & Haehnelt, 2000; Monaco et al., 2000). All of the above suggests that galaxy and quasar, both formation and evolution are closely related, not least because there is so far no evidence of quasar activity outside galaxies.

Previous work derives the black hole mass distribution from Cold Dark Matter (CDM) models (Haehnelt & Rees, 1993; Haehnelt, Natarajan & Rees, 1998) or from observations of the local non-active galaxies (Salucci et al., 1999), but then considers an essentially arbitrary exponential quasar light curve: a short active phase with luminosities close to the Eddington limit at the beginning of the black hole's life, followed by a rapid exponential decay, due to fuel exhaustion caused by the high accretion rates needed. Such a quasar light curve is only an approximation of the accretion process and does not include the physics of the accretion disk which is thought to be formed during accretion onto a supermassive black hole in quasars (e.g. Lynden-Bell 1969, Shields 1978, Rees 1984). Accretion disks have been studied in detail in many types of binary systems, where they usually show complex variability. Both observations and theory point to strong similarities in the behavior of accretion disks around Galactic black hole X-ray sources and of those

around quasars (Tanaka & Lewin, 1995; Fiore & Elvis, 1997; Czerny, Schwarzenberg-Czerny & Loska, 1999). These similarities suggest quite a different quasar light curve. Large amplitude instabilities typically appear in accretion disks, leading to repetitive outbursts as large as a factor of 10^4 on the timescale of thousand to million years depending on the disk model (Siemiginowska, Czerny and Kostyunin, 1996, Lin & Shields 1986). In the case of quasar black holes, these variations alone inevitably create a broad luminosity function (Siemiginowska & Elvis, 1997).

Unstable accretion leading to episodic outbursts has a valuable feature. In our previous paper (Siemiginowska & Elvis, 1997) quasar light curves were derived by assuming a continuous fuel supply at a low constant accretion rate of $\sim 0.04\dot{M}_{Edd}$ (where $\dot{M}_{Edd}$ is the critical accretion rate corresponding to the Eddington luminosity for a given mass). At this rate a $10^6 M_\odot$ black hole accretes some $10^{-3} M_\odot/\text{yr}$ while a $10^9 M_\odot$ black hole accretes $1 M_\odot/\text{yr}$, so a black hole would need 10^{10} years to change its mass by an order of magnitude. Consequently the problem of creating hypermassive ($10^{11} M_\odot$ Soltan 1982, Phinney 1983) quasar remnants does not occur, if the highest initial black hole mass is $10^9 M_\odot$.

In this paper we build on the Siemiginowska & Elvis (1997) work by combining the CDM derived black hole mass spectrum with the luminosity function for a particular mass derived from the time-dependent accretion disk theory. This results in an end-to-end model for quasar evolution that is completely physics based, excepting only the ratio of black hole mass to dark matter halo mass. Since this model links CDM and quasar evolution, the parameters of CDM constrain quasar evolution, and the observed quasar evolution constrains CDM parameters. We examine some of the implications of this model.

2. Model and assumptions

2.1. Accretion onto supermassive black holes

We assume that each quasar hosts a supermassive black hole surrounded by an accretion disk. The disk is the main emission component and driver of the quasar activity. We calculate a quasar light curve assuming that quasar emis-

sion is due to a non-stationary accretion disk, thus the quasar luminosity during evolution of accretion disk is calculated by integrating the disk emission at each epoch. Our time-dependent accretion disk model (Siemiginowska et al., 1996) takes into account thermal-viscous instabilities due to Hydrogen ionization, which can develop in the disk (Smak, 1982; Lin & Shields, 1986). These instabilities are observed in both cataclysmic variables and Galactic Black Hole Candidates (Smak, 1984; Tuchman, Mineshige & Shields, 1990), where they cause factor of 10^4 luminosity outbursts. Similar amplitudes are expected in the disks around super-massive black holes but the timescales are of order of hundred to hundred million years (Burdari et al 1998, Siemiginowska, Czerny & Kostyunin, 1996; Mineshige & Shields 1990). As a consequence a single source can emit within a wide luminosity range depending on the current state of the disk. Even if there is a continuous fuel supply the source will exhibit many active and quiescent states, making ‘quasar activity’ a recurrent phenomenon occurring numerous times during a galaxy’s lifetime. For example, for a black hole of $10^8 M_\odot$, 5 outbursts occur in $\sim 3.2 \times 10^6$ years (figure 1). Siemiginowska & Elvis (1997) derived the quasar luminosity function by integrating the time spent by a quasar in each luminosity state and assuming that this distribution is applied over a whole (uncoordinated) population with a range of black hole masses.

We have checked a simplifying assumption in Siemiginowska & Elvis (1997) that the distribution of time in each state is constant for all black holes masses. In figure 1 we show the light curves, due to emission from the accretion disk which exhibits thermal-viscous instabilities, for black holes with masses $10^6 M_\odot$, $10^7 M_\odot$, $10^8 M_\odot$ and $10^9 M_\odot$. The overall form of the light curves is the same, but for larger masses, outbursts are rarer than for lower masses, the high luminosity phases last much longer and the mean luminosity increases too. Figure 2 quantifies this by showing how the number of outbursts per 10^6 yr and the duration of each outburst vary with the black hole mass. Both the number of outbursts and their duration depend quadratically on the black hole mass.

Note that at $10^6 M_\odot$ an outburst occurs every century, which means that in a sample of 100 AGN we should be able to observe one transition per

year. The Piccinotti et al (1982) 2-10 keV selected sample has ~ 20 non-blazar AGNs, roughly equally divided between broad and narrow-line objects. In the ~ 20 years that these sources have been studied, one (NGC 7582) has changed from purely narrow to broad-line status, followed by a considerable change in its luminosity (Aretxaga et al., 1999). This is consistent with the expected statistics. Monitoring of large ($N \sim 1000$) emission-line selected AGN samples for a few years would strongly test this key premise of the model.

Integration of the light curves over time shows that each source spends the greater part of its lifetime (around $\sim 70\%$) in a “quiescent” state and $\sim 30\%$ in an active state ($L \geq 0.001 L_{Edd}$) with $\sim 10\%$ in a very active state ($L \geq 0.1 L_{Edd}$). These percentages are almost identical for all black hole masses, as shown in table 1. Deviations are no larger than 5% over a range of 4 orders of magnitude, which is insignificant. The fraction of time a source spends in each activity state is essentially indeed independent of its mass, as assumed by Siemiginowska & Elvis (1997).

2.2. Dark matter halos and black holes

To derive the black hole mass function, we use the Press - Schechter (1974) formalism, a simple yet quite adequate approach, which has become the basis of most formation scenarios of dark matter halos in CDM models (e.g. White & Frenk, 1991; Haehnelt & Rees, 1993). The Press - Schechter formalism predicts the mass fraction of the Universe contained in virialized structures, resulting from a gravitational hierarchical process (dark matter halos). The mass function of the dark matter halos (DMHs) can be written as:

$$n(M, z) dM = - \left(\frac{2}{\pi} \right)^{1/2} \frac{\bar{\rho}}{M} \frac{\delta_c}{\sigma^2} \frac{d\sigma}{dM} \exp \left(- \frac{\delta_c^2}{2\sigma^2} \right) dM \quad (1)$$

where $\bar{\rho}$ is the comoving density. This formula gives the number of dark matter halos of mass M formed up to a redshift z . δ_c , the critical overdensity in linear theory for spherical collapse, has been considered constant, equal to 1.69 (e.g. Sahni & Coles, 1995), although it depends weakly on Ω_0 and Λ . The linear theory rms mass density fluctuation in spheres of mass M at redshift z is

$$\sigma = \sigma_0 \frac{D_z}{D_0}$$




Fig. 1.— Quasar light curves (accretion disk luminosity vs time in 10^6 yr) due to thermal-viscous instabilities in the accretion disk around a central black hole. From top to bottom and left to right the masses of the black holes are $10^6 M_\odot$, $10^7 M_\odot$, $10^8 M_\odot$ and $10^9 M_\odot$. Accretion rate at the outer edge of the disk is equal to $0.04 \dot{M}_{Edd}$ of critical (Eddington limit) accretion rate.

where

$$D_z = (1 + 3/2\Omega_0)(1 + 3/2\Omega_0 + 5/2\Omega_0 z)^{-1}$$

is a simplified form of the linear growth factor (Peebles, 1980) that represents the growth of a perturbation in the linear regime. For the calculation of σ we used the approximation of the present rms mass density fluctuation in spheres of mass M (e.g. Fan et al., 1997):

$$\sigma_0 = \sigma_8 \left(\frac{M_8}{M} \right)^\alpha$$

where $M_8 = 1.19 \times 10^{15} \Omega_0$ is the mean mass in a sphere of radius $8h^{-1} Mpc$, $\alpha = (n + 3)/6$ where n

is the exponent of the power spectrum $P(k) \propto k^n$, and the corresponding $\sigma_8 = 0.67$. Comparing to other results (Haehnelt, Natarajan & Rees, 1998) we verified that the shape of the mass function for masses greater than $10^8 M_\odot$ is not affected by this approximation.

The DMH mass function for the standard CDM model, for $n = -1$ and for various redshifts between 5 and 0 is illustrated in figure 3a. A single cosmological model ($q_0=0.5$) will be used henceforth.

Equation 1 refers to halos harboring both elliptical and spiral galaxies. Quasars are found in both elliptical and spiral galaxies (McLeod et al., 1995; McLeod & Katris, 1995; Taylor et al., 1996;




Fig. 2.— (a) Variation of the number of outbursts (high luminosity active state) per 10⁶ yr and (b) duration of the outbursts (high luminosity state) (in 10⁶ yr) versus the black hole mass. The solid lines illustrate the linear fits.

Table 1: Fraction of time sources of different mass spent in the different activity states

$M_{BH}(M_{\odot})$	$L/L_{Edd} < 10^{-3}$	$10^{-3} < L/L_{Edd} < 10^{-1}$	$L/L_{Edd} > 10^{-1}$
10 ⁶	0.73	0.20	0.07
10 ⁷	0.75	0.15	0.10
10 ⁸	0.77	0.14	0.09
10 ⁹	0.71	0.13	0.15

Bahcall et al., 1997; Boyce et al., 1998). Narrow lined (type 2) AGN can also reside in both early type galaxies and in spirals (e.g. Salucci et al., 1999). We assume that the distinction between quasars and Seyferts is purely one of luminosity (we also include both radio-loud and radio-quiet AGN in term ‘quasar’), the latter occurring in spiral galaxies, since we consider that they are both the manifestation of the same basic phenomenon. The black hole mass function can be directly derived from the dark matter halo mass function. We assume that in each newly formed dark matter halo a black hole is “instantly” created, i.e. that all galaxies harbor a black hole. This hypothesis has been supported by several observations (e.g. Keel, 1983; Ho et al., 1997; Magorrian et al., 1998; Richstone et al., 1998; Ford et al., 1998). Haiman

& Loeb (1998) assumed that black hole formation is restricted to halos whose masses are bigger than $10^8((1+z)/10)^{-3/2}M_{\odot}$ with the mass of the black hole being a constant fraction of the dark matter halo mass: $M_{BH} = 10^{\alpha}M_{DMH}$, where $\alpha = -3.2$ for all black hole masses. We adopt this mass relation as a baseline, but examine the possibilities of varying α between -4.5 and -3.2 . The resulting black hole mass functions are shown in figure 3b. Values close to -3.2 are consistent with the $M_{BH}/M_{bulge} = 0.006$ of Magorrian et al. (1998), while values close to -4.5 are consistent with the $M_{BH}/M_{bulge} = 10^{-3.5}$, given by Wandel (1999), where M_{bulge} denotes the mass of the galactic bulge. We also consider the possibility of an α that is a function of the DMH mass, $\beta \times \log(M_{DMH})$, which is equivalent to a non-linear relation be-

tween the BH and DMH masses, $M_{BH} = M_{DMH}^{1+\beta}$. The dash-dotted line in figure 3b traces the black hole mass function for $\beta = -0.332$, a value chosen by fitting the observations, as explained in the next section.

3. Deriving the luminosity function

The luminosity function is the product of the fraction of time each source of a given mass spends in each luminosity (activity state) and their space density (Siemiginowska & Elvis, 1997). More formally, the luminosity function of a population of quasars is the result of the convolution of their mass function with the light curve, $L(t)$, of an individual source of each mass:

$$\Phi(L, z) = \int \Phi(L, z, M) n(z, M) dM, \quad (2)$$

where $n(z, M)$ is the mass function and $\Phi(L, z, M)$ is the fraction of objects that contribute to a given luminosity at a given redshift (equivalent to f_{on} in equation 3).

This is equivalent to the Haiman & Menou (2000) suggestion that the observed quasar luminosity function $dN_{obs}(z, M)/dL$, can be directly compared with the Press - Schechter DMH mass function, $dn(z, M)/dM$, to derive the luminosity of each halo of mass M :

$$\frac{d\Phi_{obs}(z, M_{BH})}{dL} \frac{dL}{dM} = f_{on} \frac{dn(z, M)}{dM}, \quad (3)$$

where f_{on} is the quasar “duty cycle” defined as the fraction of black holes that are active at a given redshift, z . Note that equation 2 is the integrated form of equation 3, if the halo mass function is replaced by the black hole mass function.

By integrating the light curves over timescales that allow many outbursts to occur, we calculated the time a source spends in the different activity states for a range of black hole masses. This is equivalent to calculating, for a certain redshift, z , the fraction of sources of a given mass, M , in a given activity state, L , $\Phi(L, z, M)$. All the above are illustrated in figure 4a, where we show the contribution of sources with black hole masses from $10^6 M_\odot$ to $10^9 M_\odot$, to the overall luminosity function, for a Press - Schechter derived black

hole mass function at a redshift of 2. The sources with black hole masses within the abovementioned range have luminosities within the observed limits ($10^{41} \text{ erg/sec} \leq L \leq 10^{48} \text{ erg/sec}$). Masses have been converted to luminosities through the light curves and α was held constant, $M_{BH} = 10^{-3.2} M_{DMH}$, for all black hole masses.

Note that in each case, a low luminosity state for a high mass source, is a high luminosity state for a lower mass source, and vice versa. Figure 4b shows that at each redshift, the slope defined by the high activity sources gives roughly the shape and slope of the luminosity function. In the case of an upper or lower mass limit, the luminosity function will follow the shape of the corresponding light curves, creating two breaks: one towards the lower and one towards the higher luminosities. Figure 5 illustrates the luminosity function of the form $\Phi(M_B) dM_B$, for different values of the parameter n (Fig. 5a) and the parameter α (Fig. 5b), including black hole masses $10^6 M_\odot \leq M_{BH} \leq 10^9 M_\odot$, after converting the luminosities into absolute magnitudes, M_B . The shape of the luminosity function and its evolution depend on these two parameters. Our discussion is focussed on the effect of varying these parameters.

3.1. Effects of changing CDM Power Spectrum Slope (n)

Here we examine the effect of changing the values of the exponent n of the CDM power spectrum. For a CDM-type spectrum $P(k) \propto k^n$, n takes values in the interval $-2 < n < -1$ on scales of $\sim 8h^{-1}$ Mpc (e.g. Fan et al., 1997). For many years, the galaxy correlation function data were consistent with $n = -1.2$ (Peebles, 1980; Davis & Peebles, 1983). However, new surveys show that the shape of the power spectrum has a more complicated form and/or that the values of n vary between small and large scales (e.g. Lin et al., 1996; Hermit et al., 1996; Le Fevre et al., 1996; Tadros et al., 1999). We do not attempt to introduce a more complex form for the power spectrum. However, we do find that varying the exponent, n produces strong effects on the quasar luminosity function. Observations show that the quasar space density increases as $\Phi(z) = \Phi_0(1+z)^3$ from the current epoch up to a turnover redshift, z_t , of 2-3, where it stops or possibly drops (e.g. Hook et al., 1998). Our models naturally produce a z_t by integrating




Fig. 3.— (a) Press - Schechter DMH mass function for $z=5$ (dotted line), 3 (solid line), 2 (dashed line) and 0 (dash-dotted line) for the cosmological model $q_0=0.5$ and for $n = -1$. (b) Black hole mass functions for $z = 3$. The solid line corresponds to $\alpha=-3.2$, the dotted line to $\alpha=-3.8$, the dashed line to $\alpha=-4.5$ and the dash-dotted line to $\alpha = -0.332 \times \log(M_{DMH})$.

over the whole redshift range the predicted luminosity function, for sources with absolute magnitudes brighter than $M_B = -26$. We find that the z_t predicted by our models is a parabolic function of n . Figure 6a shows the logarithmic normalized space density of quasars brighter than $M_B = -26$ as a function of z , for 4 different values of n (with $\alpha = -4.5$). The peak of the distribution, z_t , changes with n : the steeper the power spectrum the lower the z_t . Figure 6b shows the relation between z_t and n , which is roughly parabolic. Moreover, this relation is almost independent of the value of α making this a robust result, since α is the only physically undetermined parameter in the model.

The observed z_t is at least as high as 2.5 (Hook et al., 1998) and may be as high as ~ 4 (Miyaji et al., 2000), but quasar space density declines rapidly for $z > 4$, as suggested by both optical and radio observations (Kennefick et al., 1997; Hook et al., 1998) and theoretical models (e.g. Cavaliere & Vittorini, 1998, give a z_t around 3.5). This puts a strong constraint on n , which can only take the values between -1.6 and -1.8 (figure 6b). These values are in excellent agreement with the values

given by Blanchard et al., 1999, derived from the galaxy clusters. We restrict the rest of our study to this range of n .

3.2. Dependence of Halo to black hole mass ratio (α) on mass.

In the next two sections we discuss how the parameter α affects the luminosity function and we consider two possibilities: (1) α is constant for all masses; (2) α is related to the mass of the DMH.

3.2.1. Constant (α)

Most theoretical models have assumed for simplicity that the black hole mass is a constant fraction of the dark matter halo mass, independent of the masses involved (e.g. Haiman & Loeb, 1998; Haiman & Menou, 2000). Observationally the results of Magorrian et al (1998) suggest that this is roughly obeyed. Therefore we first examine this possibility and compare our predictions with observations, both for low and for high redshifts. Figure 7 illustrates the predicted luminosity function in the case of $\alpha = -3.2$ and $\alpha = -4.5$ for different values of n ranging from -1.6 to -1.8 , at




Fig. 4.— (a) Contribution of the different black hole mass sources to the luminosity function at $z = 2$, for black hole masses ranging from $10^6 M_\odot$ (solid line) to $10^9 M_\odot$ (dash-dotted). α was constant and equal to -3.2 for all black hole masses. (b) The slope defined by the sources at high luminosity state gives roughly the slope of the luminosity function. In the case of an upper (here $10^9 M_\odot$) or lower (here $10^6 M_\odot$) mass limit, the luminosity function will follow the shape of the corresponding light curves, creating two breaks.

$z = 3.25$. The symbols are the observed luminosity function from Warren et al. (1994).

In both cases, the slope of the predicted luminosity function is different and much smoother than the observed luminosity function and no break is present. The observed luminosity function cannot be fitted with the constant α model. For a high value of α (-3.2) the luminosity function is over-estimated at high luminosities, while for a low value ($\alpha = -4.5$) the numbers of faint quasars are under-predicted. The same tendencies are seen at all redshifts. This suggests that α cannot be constant, but may be a function of the masses involved.

3.2.2. α dependent on the M_{DMH}

There is no a priori reason that the black hole mass and the dark matter halo mass should have constant ratio, so in this section we discuss the effects of varying α . We assume that α depends on the DMH's mass with the relation $\alpha = \beta \times \log(M_{DMH})$, or, in other words, that $M_{BH} = M_{DMH}^{1+\beta}$. In order to obtain the best

value of a parameter β we fit this function to the observed luminosity function from Warren et al. (1994) at redshifts 2.6 and 3.25, shown in figure 8. We found that $\beta = -0.332^{+0.04}_{-0.016}$ (or $1+\beta = 0.668$) gives the best fit, which is the value used below.

The black hole masses are now defined from the relation: $M_{BH} = M_{DMH}^{0.668}$. This relation gives $\alpha = -3$ for $M_{BH} = 10^6 M_\odot$ and $\alpha = -4.4$ for $M_{BH} = 10^9 M_\odot$, which are in agreement with the observed values (Magorrian et al., 1998 and Wandel, 1999).

Figure 8 illustrates the predicted luminosity function for $z = 2.6$ and $z = 3.25$ for 3 values of n from -1.6 to -1.8 . The symbols show the Warren et al. (1994) data. The fit is much better than for any constant value of α . The bright end of our predicted luminosity functions at $z = 2.6$ seems to be over-predicted, when compared to observations (figure 8a). Alternatively, the observed bright end may have been under-estimated, as suggested by recent surveys, in particular the Hamburg/ESO survey (Wisotzki et al., 1996, Koehler et al, 1997 and Wisotzki, 2000). We conclude that α must be a function of the DMH in order to properly




Fig. 5.— Predicted luminosity function for $z = 2$. (a) n varies while α is kept constant, (b) α varies while n is constant.

describe the most recent observations.

Our physical model leads to changes in the luminosity function equivalent to either PLE or PDE, up to a redshift of ~ 4 , because its form is close to being a simple power law.

3.3. Comparison with observations - discussion

The predicted evolution fits the data quite well at high redshifts, e.g. $z > 2.0$. For redshifts below ~ 2 , however, the predicted luminosity function, in terms of PLE with $L = L_*(1+z)^{\gamma_p}$ evolves as $\gamma_p = 0.5$. This disagrees with the observed luminosity function issued from the following samples: AAT (Boyle et al., 1991), HES (Wisotzki et al., 2000), LBQS (Hewett et al., 1995) and the CFRS quasars (Le Fevre et al., 1995), for redshifts 0.5 to 2.0 (figure 9), that evolves much more strongly with, $\gamma_o \sim 3.5$, although the slopes of the two (observed and theoretical) luminosity functions are the same, as seen in figure 9. In this figure, the symbols show the observed luminosity function issued from the abovementioned samples while the dashed, dotted and thick solid lines illustrate the predicted luminosity function for $n = -1.8$, $n = -1.7$ and $n = -1.6$, respectively.

For many years, the best established form of the observed quasar luminosity function was a broken power law, first given for optically selected quasars with redshifts $0.3 \leq z \leq 2.2$ by Boyle et al. (1988), confirmed by La Franca & Cristiani (1997) for other optical surveys and by Page et al. (1996) for X-ray selected AGN. However, recent surveys, in several wavelengths, show that this assumption was probably due to selection effects. For $z > 2.2$ (as $\text{Ly}\alpha$ shifts from the U to the B-band) multicolor selection techniques result in biased samples and favor the redshift regions at which quasar colors make them distinguishable from stars. Giallongo and Vagnetti (1992) argued that the observed break was due to the uncertainties on the K-correction due to the dispersion in quasar optical slopes. This seems to be confirmed by ROSAT AGN data (Miyaji et al., 2000). More recent surveys, which account for this effect, such as the Hamburg/ESO Quasar Survey (which uses emission lines in addition to colors, Wisotzki et al., 1996, Koehler et al., 1997 and Wisotzki, 2000) show that the number of bright objects was underestimated and that the combined *local* luminosity function for Seyferts and quasars could be fitted with a single unbroken power law, as argued by Giallongo and Vagnetti (1992). Pei (1995) also




Fig. 6.— (a) Predicted logarithmic normalized quasar space density for $n = -1.0$ (solid line), $n = -1.3$ (dotted line), $n = -1.7$ (dashed line) and $n = -2.0$ (dash-dotted line) for $\alpha = -4.5$. (b) Variation of z_t with the power spectrum exponent, n , for $\alpha = -3.2$ (solid line), $\alpha = -4.5$ (dotted line) and $\alpha = -0.332 \times \log(M_{DMH})$. The relation is parabolic and almost independent of α . The dashed lines mark the accepted values for z_t , deduced from observations and the constraints they impose on the values of n .

suggests that a single power law of the form $L^{1/4}$ could fit the data as well as a broken power law. This single power law is represented by the thin solid line in figure 9. In terms of luminosity, galaxies with central black hole masses of the order of $10^6 M_\odot$ will be seen as Seyfert galaxies, while the higher masses will result in quasars (figure 4).

Even if the observations are altered somewhat additional physical effects are likely to be important, since we derived the mass function using only the Press - Schechter DMH mass function. For example galaxy mergers could significantly modify the assumed mass function (e.g. Haiman & Loeb, 1998; Kauffmann & Haehnelt, 2000). Mergers are rare in the early Universe but start becoming important at a redshift around 2, an epoch of massive galaxy formation (Kats et al., 1992; Kontorovich et al., 1992). Only mergers that result in the formation of a single object will cause a decrease of the space density of the black holes. The black holes, due to the dynamical friction, will soon be found in the center of the newly formed object and merge (e.g. Binney & Tremaine, 1994), thus influencing the quasar space density. If this is the

primary cause of the low z disagreement of our model with observation then the difference of the observed to predicted evolution rates, $\gamma_o - \gamma_p$, provides a crude estimate of the merger rate, γ_m , for redshifts between 0.5 and 2, i.e. $\gamma_m = 3$. Hence the merger rate can be expressed as a function of the redshift: $\sim (1+z)^3$. Interestingly this value is in agreement with several other results based on observations: Zepf & Koo (1989) studied a complete sample of 20 close pairs of faint ($B \leq 22$) galaxies and found that the frequency of interacting galaxies increase as $(1+z)^{4.0 \pm 2.5}$. Burky et al. (1994) used four fields observed with the Hubble Space Telescope and found that the galaxy merger rate increases with redshift as $(1+z)^{2.5 \pm 0.5}$. Patton et al. (1997) used a sample of 545 field galaxies and estimated that the merger rate changes with redshift as $(1+z)^{2.8 \pm 0.9}$.

Note that less extreme galaxy interactions, where the objects remain independent entities afterwards will not influence our results, if we assume that whatever happens in the vicinity of an object will not influence its accretion rate and therefore its light curve. In reality, such interac-




Fig. 8.— Predicted luminosity function for $z = 2.6$ and 3.25 , for $n = -1.6$ (solid line), -1.7 (dotted line) and -1.8 (dashed line). Symbols show the observed luminosity function of Warren et al. (1994).

tions are likely to affect the accretion rate and will be taken into account in the future work.

The disagreement between the observed and the predicted luminosity function at $z \leq 2$ may also be due to the role of obscured (type II) AGN. The estimates of their number density are not yet accurate. Recent work (Salucci et al., 1999) estimates the local mass function of the dormant black holes within elliptical galaxies, assuming that the nuclear activity is a single short event. Salucci et al., however, do not take merger effects into account, introducing uncertainties in the calculated massive dark object mass function.

We used a black hole mass distribution with black holes ranging from $\sim 10^6 M_\odot$ to $10^9 M_\odot$. According to our model, accretion onto such black holes can produce luminosities within the observed range $10^{41} \leq L \leq 10^{48}$ erg/sec ($10^{-4} \leq L/L_{Edd} \leq 1$). If we allow for a higher black hole mass cut-off at $\sim 10^7 M_\odot$ we could obtain the break in the luminosity function. However, there is evidence for black holes with lower masses (e.g. Reid et al., 1999, Sag A* $\sim 2.6 \times 10^6 M_\odot$). Therefore a break cannot be imposing a simple cut-off mass.

Our model gives a consistent physical connection between high and low luminosity sources and

does not predict a break in the luminosity function for the black hole mass distribution within the observed mass range.

The model presented here can be extended and used for the study of the UV-background and the re-ionization history of the Universe (Siemiginowska et al., 2000, in preparation). Issues such as whether quasars are entirely responsible for the re-ionization of the Universe or whether Population III stars are required (see Madau 1999), the redshift at which it occurred, whether it is homogeneous or inhomogeneous or its effects on the primordial power spectrum, can be examined. The results are likely to be different from previous attempts because the emission spectrum derived from a non-stationary accretion disk is different from the one predicted by a stationary model at the UV wavelength range (Siemiginowska et al., 1997).

4. Conclusions

In this paper we presented a theoretical model for calculating the quasar luminosity function. This model, unlike previous ones, takes into account the physical processes occurring in the accretion disk around a black hole. Black holes are




Fig. 7.— Predicted luminosity function at $z = 3.25$ for different values of n for constant α . Upper set of curves: $\alpha = -3.2$, lower set of curves: $\alpha = -4.5$. The symbols show the observed luminosity function from Warren et al (1994).

assumed to form instantly within all dark matter halos, with masses related to the halo mass as: $M_{BH} = M_{DMH}^{0.668}$.

We first restricted the range for the two free parameters of the model: the power spectrum exponent, n and the ratio of black hole mass to DMH mass, α . For any reasonable α value, $(-4.5 < \alpha < -3.2)$, n has to be within the narrow range -1.6 to -1.8 in order to match the observed redshift of the peak of quasar evolution. We compared the predicted luminosity function to the data and found that a non-linear relation between the BH and DMH masses ($M_{BH} = M_{DMH}^{1+\beta}$) gives a much better approximation than a linear relation ($M_{BH} = 10^\alpha M_{DMH}$). Our model predicts no break in the luminosity function, assuming quasars and Seyfert galaxies to be objects of the same kind but of different luminosities. Observationally this model is then equivalent to both PLE and PDE, but physically has quite different implications.

No assumptions have been made on the merger processes, although they impact the quasar evolu-

tion. Mergers that result in the black hole fusion (and consequently in a decrease of the black hole space density) will also influence the luminosity function at redshifts lower than ~ 2 , when mergers become important. We estimated that the merger rate should depend on redshift, $(1+z)^3$, consistent with direct measurements.

Built into our model is the idea that quasar-like activity is a recurrent phenomenon. Practically all galaxies go through that state many times during their lives. Recent observations support the idea that black holes reside in all galaxies.

This work was supported by NASA grants NAG5-3391, NAG5-4808, NAG5-6078, and NAS8-39073.

REFERENCES

- Aretxaga, I., Joguet, B., Kunth, D., Melnik, J., Terlevich, R. J., 1999, *ApJ*, 519, L123
- Bahcall, J. N., Kirhakos, F., Saxe, D. H., Schneider, D. P., 1997, *ApJ*, 479, 642
- Binney, J. & Tremaine, S., 1994, *Galactic Dynamics*, Princeton University Press, 1994, p.422
- Blanchard, A., Sadat, R., Bartlett, J. G. & Le Dour, M., 1999, submitted to *A&A*, preprint astro-ph/9908037
- Boyce, P. J., Disney, M. J., Blades, J. C., Boksenberg, A., Crane, P., Deharveng, J. M., Macchetto, F. D., Mackay, C. D., Sparks, W. B., 1998, *MNRAS*, 298, 121
- Boyle, B.J., Shanks, T. & Peterson, B.A., 1988, *MNRAS*, 235, 935
- Boyle, B.J., Jones, L. R. & Shanks, T., 1991, *MNRAS*, 251, 482
- Burderi, L., King, A. R. & Szuszkiewicz E., 1998, *ApJ* 509, 85.
- Burkey, J. M., Keel, W. C., Windhorst, R. A. & Franklin, B. A., 1994, *ApJ*, 429, L13
- Cavaliere, A. & Padovani, P., 1988, *ApJ*, 333, L33
- Cavaliere, A. & Padovani, P., 1989, *ApJ*, 340, L5
- Cavaliere, A. & Vittorini, V., 1998, *The Young Universe: Galaxy Formation and Evolution at Intermediate and High Redshift*. Edited by S. D'Odorico, A. Fontana, and E. Giallongo. ASP Conference Series; Vol. 146; 1998, p.26

- Colbert, E. J. M. & Mushotzky, R. F., 1999, *ApJ*, 519, 89
- Czerny, B., Schwarzenberg-Czerny, A. & Loska, Z., 1999, *MNRAS*, 303, 148
- Davis, M. & Peebles, P. J. E., 1983, *ApJ*, 267, 465
- Efstathiou, & Rees, M. J., 1988, *MNRAS*, 230, 5p
- Fan, X., Bahcall, N. A. & Cen, R., 1997, *ApJ*, 490, L123
- Fiore, F. & Elvis, M., 1997, *Adv. Space. Res.*, 12(1), 85
- Ford, H. C., Tsvetanov, Z. I., Ferrarese, L., Jaffe, W., 1998, in Sofue Y., ed., *Proc. IAU Symp.* 184, *The Central Regions of the Galaxy and Galaxies*, Kluwer, Dordrecht, p.377
- Giallongo, E. & Vagnetti, F., 1992, *ApJ*, 396, 411
- Haehnelt, M. G. & Rees, M. J., 1993, *MNRAS*, 263, 168
- Haehnelt, M. G., Natarajan, P. & Rees, M. J., 1998, *MNRAS*, 300, 817
- Haiman, Z. & Loeb, A., 1998, *ApJ*, 503, 505
- Haiman, Z. & Menou, K., 2000, *ApJ*, 531, 42
- Hermit, S., Santiago, B.X., Lahav, O., Strauss, M.A., Davis, M., Dressler, A. & Huchra, J.P., 1996, *MNRAS*, 283, 709
- Hewett, P. C., Foltz, C. B. & Chaffee, F. H., 1995, *AJS*, 109, 1498
- Ho, L. C., Filippenko, A. V. & Sargent, W. L. W., 1997, *ApJ*, 487, 568
- Hook, I. M., Shaver, P. A. & McMahon, R. G., 1998, *The Young Universe: Galaxy Formation and Evolution at Intermediate and High Redshift*. Edited by S. D'Odorico, A. Fontana, and E. Giallongo. ASP Conference Series; Vol. 146; 1998, p.17
- Kats, A. V., Kontorovich, V. M. & Krivitsky, D. S., 1992, *A&A Trans.*, 3, 53
- Kauffmann G. & Haehnelt, M., 2000, *MNRAS*, 311, 576
- Keel, W. C., 1983, *ApJ*, 269, 466
- Kennefick, J. D., Osmer, P. S., Hall, P. B. & Green, R. F., 1997, *AJ*, 114, 2269
- Kohler, T., Groote, D., Reimers, D. & Wisotzki, L., 1997, *A&A*, 325, 502
- Kontorovich, V. M., Kats, A. V. & Krivitsky, D. S., 1992, *Sov. Phys. JETP Lett.*, 55, 1
- Krolik, 1999, *Active Galactic Nuclei*. Princeton University Press.
- La Franca, F. & Cristiani, S., 1997, *AJ*, 113, 1517
- Le Fevre, O., Crampton, D., Lilly, S.J., Hammer, F. & Tresse, L., 1995, *ApJ*, 455, 60
- Le Fevre, O., Hudon, D., Lilly, S.J., Crampton, D., Hammer, F. & Tresse, L., 1996, *ApJ*, 461, 534
- Lin, H., Kirshner, R. P., Shectman, S. A., Landy, S. D., Oemler, A., Tucker, D. L. & Schechter, P. L., 1996, *ApJ*, 471, 617
- Lin, D. N. C. & Shields, G. A., 1986, *ApJ*, 305, 28
- Lynden Bell, D., 1969, *Nature*, 223, 690
- Madau, P. 1999, in: *After the Dark Ages: When Galaxies were Young (the Universe at $2 < z < 5$)*. 9th Annual October Astrophysics Conference in Maryland held 12-14 October, 1998. College Park, Maryland. Edited by S. Holt and E. Smith. American Institute of Physics Press, p. 299
- Magorrian, J., Tremaine, S., Richstone, D., Bender, R., Bower, G., Dressler, A., Faber, S. M., Gebhardt, K., Green, R., Grillmair, C., Kormendy, J., Lauer, T., 1998, *AJ*, 115, 2285
- Mathez, 1976, *A&A*, 53, 15
- McLeod, K. K., Rieke, G. H., Storrie-Lombardi, L. J., 1999, *ApJ*, 511, L67
- McLeod, K. K., Katris, K., 1995, *PASP*, 107, 91
- Mineshige, S. & Shields, G. A. 1990, *ApJ* 351, 47
- Miyaji, T., Hasinger, G. & Schmidt, M., 2000, *A&A*, 353,
- Monaco, P., Salucci, P. & Danese, L., 2000, *MNRAS*, 311, 278
- Page, M. J., Carrera, F. J., Hasinger, G., Mason, K. O., McMahon, R., Mittaz, J. P. D., Barcons, X., Carballo, R., Gonzalez-Serrano, I. & Perez-Fournon, I., 1996, *MNRAS*, 281, 579
- Patton, D. R., Pritchet, C. J., Yee, H. K. C., Ellingson, E. & Carlberg, R. G., 1997, *ApJ*, 475, 29
- Peebles, P. J. E., 1980, *The Large Scale Structure of the Universe*. Princeton University Press.

- Pei, Y.C., 1995, ApJ, 438, 623
- Phinney, E.S., 1983, PhD thesis, Univ. Cambridge, England
- Piccinotti, G., Mushotzky, R. F., Boldt, E. A., Holt, S. S., Marshall, F. E., Serlemitsos, P. J. & Shafer, R. A., 1982, ApJ, 253, 485
- Press, W. H & Schechter, P., 1974, ApJ, 193, 437
- Rees, M.J., 1984, ARA&A, 22, 471
- Reid, M.J., Readhead, A.C.S., Vermeulen, R.N. & Treuhaft, R.N., 1999, in *The Central Parsecs, Galactic Center Workshop*, Tucson, Sep. 7-11, 1998. To be published in the ASP Conf. Ser.
- Richstone, D., Ajhar, E. A., Bender, R., Bower, G., Dressler, A., Faber, S. M., Filippenko, A. V., Gebhardt, K. et al., 1998, Nature Suppl., 395,14
- Sahni V. & Coles P.C., 1995, Physics Reports, 262, 135
- Salucci, P., Szuszkiewicz, E., Monaco, P. & Danese, L., 1999, MNRAS, 307, 637
- Schmidt, M., 1968, ApJ, 151, 393
- Siemiginowska, A., Czerny, B. & Kostyunin, V., 1996, ApJ, 458, 491
- Siemiginowska, A. & Elvis, M., 1997, ApJ, 482, L9-L12
- Smak, J.I., 1982, Acta Astron, 32, 199
- Smak, J.I. 1984, PASJ, 96, 5
- Soltan, A., 1982, MNRAS, 200,155
- Tanaka, Y. & Lewin, W. H. G., 1995, in *X-ray Binaries*, ed. W. H. G. Lewin, J. van Paradijs & E. P. J van den Heuvel (Cambridge: Cambridge University Press). 126
- Tadros, H., Ballinger, W.E., Taylor, A.N., Heavens, A.F., Efstathiou, G., Saunders, W., Frenk, C.S., Keeble, O., McMahon, R., Maddox, S.J., Oliver, S., Rowan-Robinson, M., Sutherland W.J & White, S.D.M., 1999, MNRAS, 305, 527
- Taylor, G. L., Dunlop, J. S., Hughes, D. H., Robson, E. I., 1996, MNRAS, 283, 930
- Tuchman, Y, Mineshige, S. & Wheeler, J.C. 1990, ApJ 359, 164
- Wandel, A., 1999, ApJ, 519, L39-L42
- Warren, S.J., Hewett, P.C & Osmer, P.S., 1994, ApJ, 421, 412
- White, S.D.M. & Frenk, C. S., 1991, ApJ, 379, 52
- Wisotzki, L., Kohler, T., Groote, D. & Reimers, D., 1996, A&ASS, 115,227
- Wisotzki, L., 2000, A&A, 353, 853
- Yi, I., 1996, ApJ, 473, 645
- Zepf, J. E. R. & Koo, D. C., 1989, ApJ, 337, 34




Fig. 9.— The predicted quasars luminosity function for $-1.8 < n < -1.6$ at (a) $z = 0.5$, (b) $z = 1.0$, (c) $z = 1.5$ and (d) $z = 2.0$ (thick lines). The symbols represent the observed luminosity function for the samples AAT (Boyle et al., 1991), LBQS (Hewett et al., 1995), HES (Wisotzki et al., 2000) and CFRS (Le Fevre et al., 1995) (Wisotzki, private communication). The thin solid line represents the single power law LF of Pei (1995).